

## Malocclusion and Premature Teeth Loss: its Prevalence and Association among Yemeni School children in Sana'a

Tharwa Ali Mokred AL-Awadi<sup>1</sup>, Khaled A AL-Haddad<sup>1</sup>, Ameen Abdullah Yahya Al-Akwa<sup>1</sup>, Mohammed Mohammed Ali Al-Najhi<sup>2</sup>, Hassan Abdulwahab Al-Shamhy<sup>3,4\*</sup>, Mohammed A Al-labani<sup>1</sup> and Omar Ahmed Esmail Al-dossary<sup>3</sup>

<sup>1</sup>Orthodontics, Pedodontics and Prevention Department Faculty of Dentistry, Sana'a University, Yemen

<sup>2</sup>Orthodontics, Pedodontics and Prevention Department Faculty of Dentistry, Genius University for Sciences and Technology, Dhamar city, Republic of Yemen

<sup>3</sup>Department of Basic Sciences, Faculty of Dentistry, Sana'a University, Republic of Yemen

<sup>4</sup>Medical Microbiology department, Faculty of Medicine, Genius University for Sciences and Technology, Dhamar city, Republic of Yemen

\***Corresponding author:** Hassan Abdulwahab Al-Shamhy, Faculty of Medicine and Health Sciences, Sana'a University

**Citation:** AL-Awadi TAM, AL-Haddad KA, Al-Akwa AA, Al-Najhi MMA, Al-Shamhy HA, et al. (2021) Malocclusion and Premature Teeth Loss: its Prevalence and Association among Yemeni School children in Sana'a. *J Oral Med and Dent Res.* 2(2):1-08.

**Received:** November 16, 2021 | **Published:** November 30, 2021

**Copyright**© 2021 by AL-Awadi TAM. All rights reserved. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

**Objectives:** This study was conducted in Sana'a city, Yemen to detect the prevalence of early loss of primary teeth and its consequence on malocclusion.

**Materials and methods:** This cross-sectional observational study included 1079 school children aged 7-12 years, and for the detection of the following characteristics, the samples were clinically examined: Angle's classification of malocclusion, anterior open-bite, overjet, lateral open-bite, overbite midline shift and cross-bite. Also, the early loss was classified according to the chronological age of an eruption of the permanent teeth proposed by Kronfeld.

**Results:** The prevalence of malocclusion among school children was 81.1% and the normal first-class molar relationship was found only in 18.9%, while the class I malocclusion included the highest rate of the sample 70.4%, followed by the class II with a 9.5% relationship, the class III included only 1.1%.

There was an increased association between early tooth loss and the development of malocclusion as this association was 0.2 times in Class I ( $p < 0.001$ ), increased in Class II to 2.2 times (0.01), and then increased to 3.7 times in Class III ( $p = 0.02$ ).

**Conclusion:** This result underscores the significance of raising awareness regarding this dilemma and the requirement to focus further efforts towards prevention of early tooth loss and interference to maintain dental health, thus recovering chewing functions and aesthetics for individuals and the entire population.

### **Keywords**

Malocclusion; Premature loss; Prevalence; Primary teeth

## **Introduction**

Investigation and survey of dental health problems in Yemen are in spite of everything fairly small and limited, even though there were studies that dealt with the problems of dental caries, the prevalence and pattern of third molar impaction in adults and children, periodontal infections, and causes of extraction of permanent teeth [1-8], however no exploration converge the prevalence of premature loss of primary teeth and its effect on malocclusion. Early loss of primary teeth is a worldwide problem. Premature loss is described as the loss of a deciduous tooth prior to the age of normal exfoliation [9]. The most common cause of early tooth loss is tooth decay, trauma, periodontal disease and early root resorption [7,8,10-12]. Early loss of primary teeth leads to early or late eruption of subsequent teeth [12]. Pediatric patients may suffer various effects, for instance anti-extrusion, dental rotation, dental crowding, craniofacial development disorders, development of harmful habits, especially post-impact tooth and reduction in dental arch length [12]. The association between the early tooth loss prevalence; and ethnicity and the environmental factors has been confirmed by many studies [9-12]. Malocclusion is a general dental health problem, its psychological and functional impact is significant, and its prevalence among children is high [13]. The reasons of malocclusion are environmental or genetic and/or a mix of equally factors, alongside with a variety of local factors for instance harmful oral habits, the shape of the teeth, dental abnormalities and teeth location of growth, can cause malocclusion [7,8,13, 14]. Regarding the early loss of deciduous teeth, it was earlier mentioned so as to the early loss of primary teeth can have an effect on the time of natural eruption of permanent successors by whichever inhibiting or speed up their eruption [15]. It is reflected on a qualifying cause for occlusal and location variances in mixed and permanent teeth [16]. It is normally recognized that the early loss of primary teeth, particularly the molar, might be in the lead to a deficiency from outer space, malocclusion and midline differences in permanent teeth [17]. Additionally, the early loss of the primary teeth reduces the arch length necessitated for the following teeth, and as a result, it pre-eliminates impaction crowding, and rotation of permanent teeth [18]. The present study was designed to ascertain the rate of the premature loss of primary teeth and its effect on malocclusion in Sana'a city, Yemen.

## **Materials and Methods**

Among Yemeni school children in primary schools (public and private) in Sana'a, Yemen, this cross-sectional descriptive study was conducted to determine the rate of malocclusion related with early tooth loss. From the selected Sana'a schools, a total of 1079 boys and girls aged 7-12 years were randomly

selected and screened. Selected schools are located in different areas of the city to avoid having children from the same area (Table 1).

The criteria factors for every malocclusion measurement described by Angle, [19] were divided malocclusion into various groups that were recognized on the occlusal relations of the first molar. Using the standard method [20], data were collected by one examiner, the first researcher (Tharwa). With a simple examination under sufficient light, sometimes if natural light was insufficient, artificial light from a torch was used a clinical examination was done for each child. The children who were selected and refused to participate were replaced by new children.

## Ethical Approval

The written consent in all cases was obtained. Approval was obtained from the participants prior to including in the study. Ethical approval was obtained from the Medical Research and Ethics Committee of the Faculty of Medicine and Health Sciences, Sana'a University with reference number (852) on 11/11/2018.

## Data Analysis

Data were investigated and obtainable using tables where percentage (%) was used to describe qualitative variables. The odds ratio (OR), 95% confidence interval, chi-square with Yate correction and Fisher's tests were used to illustrate the significance of the association between early loss of primary teeth and incidence of malocclusion at a significance level less than 0.05 (P) by SPSS software (IBM Corp. Released 2012).

## Results

Table 1 shows the distribution according to age, gender, type of school and school district for the 1079 selected children who participated in this study. The study included almost the same number for age groups where the number of children aged 7-9 years was 48.8% and 10-12 years old was 51.2%, as well as in terms of gender, the number of boys was 50.6% and girls were 49.4%. Regarding school types, 70.5% of the children selected were from public schools and only 29.5% were selected from private schools. With regard to the districts, most of the children were from Al-Thawrah district (62%) (large population density), while 15.8% were from Old Sana'a and 22.2% were from Al-Safiah. The outcomes in Table 2 show that the totality malocclusion was 81.1%, and the Class I malocclusion included the highest percentage of the sample 70.4%, finding 73.2% for the age group of 7-9 years higher than that of the age group of 10-12 years of 67.8%. This difference was statistically significant ( $\chi^2 = 6.24$ ,  $P < 0.05$ ). The second class was present at 9.5%, while the third class contained only 1.1% of the entire sample. Consistent with gender and school type, a statistically non-significant difference ( $P > 0.05$ ) was found in the incidence of malocclusion. The current study was planned to determine the statistical relationship between early tooth loss and the development of malocclusion. There was an increased association between early tooth loss and the development of malocclusion as this association was 0.2 times in Class I ( $p < 0.001$ ), increased in Class II to 2.2 times (0.01), and then increased to 3.7 times in Class III ( $p = 0.02$ ) (Table 3).

Variables		Frequency	%
Age	7-9yrs	527	48.8
	10-12yrs	552	51.2
Gender	Boys	546	50.6
	Girls	533	49.4
School type	Government	761	70.5
	Private	318	29.5
	Al- Thawrah	669	62
Districts	Old Sana'a	170	15.8
	Al- Safiah	240	22.2
Mean age $\pm$ SD= 9.5 $\pm$ 1.7			

**Table 1:** The distribution of schoolchildren participants in the study according to age, gender, school type and district (n=1079)

Angle classification	Age (year)				Gender				School type				Total N=1079	
	7-9 yrs N=527		10-12 yrs N=552		Boys N=546		Girls N=533		Government N=761		Private N=318			
	freq.	%	freq.	%	freq.	%	freq.	%	freq.	%	freq.	%	freq.	%
Class I normal	83	15.8	121	21.9	103	18.9	101	18.9	141	18.5	63	19.8	204	18.9
Total malocclusion	444	84.3	431	78.1	443	81.1	432	81.1	620	81.5	255	80.2	875	81.1
Class I	386	73.2	374	67.8	384	70.3	376	70.5	540	71.0	220	69.2	760	70.4
Class II	52	9.9	51	9.2	53	9.7	50	9.4	70	9.2	33	10.4	103	9.5
Class III	6	1.1	6	1.1	6	1.1	6	1.1	10	1.3	2	0.6	12	1.1
P-value	0.0085*				0.998				0.666					
$\chi^2$	6.24				0.035				1.57					

**Table 2:** Prevalence of Angle classification according to age, gender and school type.

Angle classification	With no premature loss (n = 565)		With premature loss (n = 310)		Odds ratio	95% CI	P
	No	%	No	%			
Class I normal	165	29.2	39	12.6	0.3	0.2-0.57	<0.001
Class I Malocclusion	512	90.6	248	80	0.3	0.2- 0.5	<0.001
Class II Malocclusion	49	8.7	54	17.4	2.2	1.5-3.3	0.01
Class III Malocclusion	4	0.88	8	2.3	3.7	1.1-12.4	0.02
Mid-line shift	237	41.9	156	50.3	1.3	1.0-1.7	0.05
Cross-bite Unilateral	109	19.3	75	24.2	1.3	0.9-1.8	0.08
Cross-bite Bilateral	23	4.1	25	8.1	2.1	1.2-3.7	0.01
Anterior open bite	65	11.5	28	9	0.7	0.4-1.2	0.25

Unilateral posterior open bite	11	1.9	8	2.6	1.3	0.5-3.3	0.5
Bilateral posterior open bite	11	1.9	8	2.6	1.3	0.5-3.3	0.5
Abnormal Over jet	323	57.2	189	61	1.2	0.88-1.5	0.2
Abnormal Over-bite	311	55	177	57.1	1.2	0.88-1.5	0.26

**Table 3:** Relationship between malocclusion with premature loss and without premature loss of the primary teeth.

With regard to over jet, the current study accounted that 323 (57.2%) of children who did not lose a premature primary tooth had an abnormal over jet compared to 189 (61%) of children who lost premature primary tooth. Concerning abnormal Over-bite, the current study found that 311 (55%) children of who did not lose their premature primary teeth had an abnormal Over-bite matched up to 177 (57.1%) of children who lost premature primary teeth.

## Discussion

Loss of premature primary teeth is an apprehension not just due to function loss on the contrary also due to the increased likelihood of other teeth erosion [18]. Prediction of subsequent primary tooth loss may be beneficial in formative treatment [21]. Furthermore, the loss of premature deciduous teeth affects the progress of the natural occlusion and generates an increased require for orthodontics treatment [21]. Primary early loss of teeth has been noticed in explore studies in numerous regions in the globe [7,8,22–25]. In the current study, 1079 children were examined, they aged 2-12 years, 318 of whom were in private schools and 761 were in public schools. The selected children were from the city of Sana'a, Yemen. Of the 1079 children, 875 (81.1%) had at least one premature loss that was either deciduous or molar canines, while 51% of children studied in Saudi Arabia [26] and 24.9% of children studied in Brazil [9] had early tooth loss. Compared to Saudi results, Danish and Brazilian, the children in the current investigation experienced significantly higher rates of premature loss of deciduous teeth. The dissimilarity could be connected to the higher mean of decayed teeth, which was found to be 4.0. It may also be for the reason that parents do not care about the primary teeth because they have the idea that the deciduous teeth will be replaced. Furthermore, in spite of the high rate of early tooth loss, upon inspection there was only two children wore space maintainers.

It has been stated and confirmed that early loss of primary teeth can affect the natural eruption time of permanent successors by either inhibiting or accelerating their appearance [23].

The etiology of malocclusion is to some extent controversial, conversely, it is simply multi-factorial, with effects being equally hereditary [27] and ecological [28]. There are three normally expected etiological factors for malocclusion: structural factors - the shape, size and virtual positions of the upper and lower jaws. Variants can be produced by environmental or behavioral factors such as masticatory muscles, nocturnal mouth breathing, cleft lip and cleft palate. The influence of muscle factors - and the function and shape of the muscles that frame the teeth - may be the cause of malocclusion. This can be influenced by habits for instance pacifier, finger-sucking, tongue-thrusting and nail-biting [29]. Dental reasons - the size of the teeth relative to the jaw, early loss of teeth may result in spacing or median migration causing crowding, abnormal eruption course or timing, extra teeth (hypernumerous), or too few teeth

(hypodontia). There is no particular cause of malocclusion, and whilst planning orthodontic therapy it is regularly helpful to be concerned about the above factors and their effect on malocclusion. These influenced by oral habits and pressure which leads to malocclusion [26,29-31].

The current study was planned to determine the statistical association between premature tooth loss and the occurrence of malocclusion. There was an increased association between premature tooth loss and the development of malocclusion, where this association was 0.2 times in the first class ( $<0.001$ ), and increased in the second class to 2.2 times (0.01), and then increased to 3.7 times in the third class ( $p = 0.02$ ) (Table 3). The outcomes of the present study definite the importance of premature tooth loss as a major factor for the development of malocclusion in Sana'a city as reported elsewhere [23,26,31]. Additionally, the results of the current study confirmed that premature loss of primary teeth is the most common local factor that leads to malocclusion due to its interfering with the harmony of adult teeth, leading to crowding initiated by migration of adjacent teeth [31].

Regarding the over jet, the current study reported that 323 (57.2%) of children who did not lose a primary tooth prematurely had an abnormal over jet compared with 189 (61%) of children who lost an premature primary tooth. This is supported by the results of Proffit et al. [32] who showed that, between 8-11 years, 45.2% of children had a slight increase in the over jet also by the findings of Al-Shahrani et al. [26] where they showed that, between 9-11 years, 53 (62.3%) children who did not lose a primary tooth prematurely had a slight increase in the over jet compared to 41 (57.7%) of the premature primary tooth loss who had mild impairment increased over jet [26]. Overbite is a vertical measurement of the degree of overlap between the upper and lower incisors. Regarding abnormal overbite, the current study found that 311 (55%) of children who did not lose their premature primary teeth had abnormal overbite compared to 177 (57.1%) of children who lost premature primary teeth. Tauscher et al [33] showed that increased overbite and over jet were the most common malocclusions in the early mixed dentition period. As a result, the extensive variety of orthodontic-related traits identified in the current study underscores the necessity for orthodontic screening at or before 9 years of age. Additionally, this study put emphasis on the importance of early detection of premature primary tooth loss to avoid future malocclusion.

Therefore, the early loss of deciduous teeth affects the development of the natural occlusion and generates an increased require for orthodontic intervention. Finally, due to the adverse effects of early loss of primary teeth, it is necessary to augment the mouth Health awareness through the implementation of school dental health programs to bring up to date children and their parents of the harmful effects of early loss of primary teeth. The value of primary teeth, children and their parents should be made aware of them to care for their preservation. Also, children with early loss of primary teeth should be educated to preserve space if required. Parents of such children also should be recommended to bring their children to the dental hospital for dental procedures.

## Conclusion

There was a high rate of malocclusion, with Class I malocclusions being the most common. There were higher rates of early loss of deciduous teeth with a significant association between malocclusion and early tooth loss. This result underscores the significance of raising awareness about this problem and the prerequisite to focus more efforts towards prevention of early tooth loss and intervention to maintain

dental health, thus improve chewing functions and aesthetics for individuals and the entire population. Additionally, our results underscore the magnitude of early recognition of early loss of primary teeth to prevent malocclusion.

## Acknowledgments

The authors extend their thanks and appreciation to Genius University of Science and Technology, Dhamar City, Republic of Yemen, which supported this work, in particular Dr. Mohammed Mohammed Ali Al-Najhi, the generous scholar who usually supports medical education and research in Yemen.

## References

1. Shoga Al-Deen, Al-Ankoshy AAM, Al-Najhi MMA, Al-Kabsia TA, AL-Haddad KA, et al. (2021) Porphyromonas gingivalis: biofilm formation, antimicrobial susceptibility of isolates from cases of Localized Aggressive Periodontitis (LAP). Universal J Pharmaceutical Res. 6(4):1-7. Alhasani AH, Ishag RA, Yahya Al-Akwa AAY, Al Shamahy HA, Al-labani MA. (2020) Association between the Streptococcus mutans biofilm formation and dental caries experience and antibiotics resistance in adult females. Universal J Pharm Res. 5(6):1-3.
2. Bamashmoos KAO, AH Alhasani, AAY Al-Akwa, AQM Q Zabara, et al.. PREVALENCE OF PREMATURE LOSS OF PRIMARY TEETH AT THE AGE OF 6-10 YEARS IN SANA'A CITY, YEMEN. Universal J Pharmaceutical Res. 5(4):1-4.
3. AL-Haddad KA, Ali Al-Najhi MM, Al-Akwa AAY, Al-Shamahy HA, Al-Sharani AA, et al. (2021) Antimicrobial susceptibility of Aggregatibacter actinomycetemcomitans isolated from Localized Aggressive Periodontitis (LAP) Cases. J Dent Ora Heal Ad Re. 2007:103.
4. Mutaher NJA, AL-Haddad KA, Al-Akwa AAY, Al-labani MA, Al-Shamahy HA, et al. (2020) Prevalence and causes of traumatic dental injuries to anterior teeth among primary school children in Sana'a city, Yemen. Universal J Pharm Res. 5(3):38-43.
5. Alhadi YAH Rassem, HA Al-Shamahy, KM Al-Ghaffari. (2019) "Causes for extraction of permanent teeth in general dental practices in Yemen". Universal J Pharmaceutical Res. 4(2):1-6.
6. Yehia LAB, K A AL-Haddad, M.A Al-labani, HA Al-Shamahy, HM. Shaga-aldeen. (2020) "Occlusal characteristics of the primary dentition among a sample of Yemeni pre-school children". Universal J Pharmaceutical Res. 5(1):359-362.
7. AL-Awadi TAM, KA AL-Haddad, MA Al-labani, HA Al-Shamahy, HM Shaga-aldeen. (2020) Prevalence of malocclusion among Yemeni children of primary schools. Universal J Pharmaceutical Res. 5(1):1-6.
8. Cavalcante A, Alencar C, Medeirosbezerra P, Granvillegarcia A. (2008) Prevalence of early loss of primary molars in school children in Campina Grande, Brazil. Pak Oral Dent J. 28: 113-16.
9. Borum Mk, Andreason JO. (1998) Sequelae of trauma to primary maxillary incisors. I. Complication in the primary dentition. Endod Dent Traumatol. 14:31-44.
10. Cardoso L, Zembruski C, Femandes DSC, Boff I, Pessin V. (2005) Evaluation of prevalence of malocclusion un relation to premature loss of primary teeth. Pesq Bras Odontoped Clin Integ. 5:17-22.
11. Heilborn J, Kuchler E, Fidalgo T, Antunes L, Costa M. (2011) Early primary tooth loss: prevalence, consequence and treatment. Int J Dent Recife. 10(3):10-13.
12. Mtaya M, Brudvik P, Astrom AN. (2009) Prevalence of malocclusion and its relationship with sociodemographic factors, dental caries, and oral hygiene in 12 to 14 years old Tanzanian schoolchildren. European Journal of Orthodontics 2009; 31(5): 467-76.
13. Shivakumar KM, Chandu GN, Subba RVV, Shafiulla MD. (2009) Prevalence of malocclusion and orthodontic treatment needs among middle and high school children of Davangere city, India by using Dental Aesthetic Index. J Indian Soc Pedod Prev Dent. 27: 211-18.
14. Leite-Cavalcanti A, Menezes SA, Granville-Garcia A.F, Correia-Fontes LB. (2008) Prevalence of early loss of

- primary molars: Study retrospective. *Acta Sci Health Sci.* 30(2):139-43
15. Baskaradoss J.K, Geevarghese A, Roger C, Thaliath A. (2013) Prevalence of malocclusion and its relationship with caries among schoolchildren aged 11-15 years in Southern India. *Korean J Orthod.* 43(1):35-41.
  16. Alamoudi N. (1999) The Prevalence of crowding, attrition, midline discrepancies and premature tooth loss in the primary dentition of children in Jeddah, Saudi Arabia. *J Clin Pediatr Dent.* 24:53-8.
  17. Lin YT, Lin WH, Lin YTJ. (2007) Immediate and six-month space changes after premature loss of a primary maxillary first molar. *J Am Dent Assoc.* 138:362-8.
  18. Angle EH. (1899) Classification of malocclusion. *Dent Cosmos.* 41:248-64.
  19. Thilander B, Pena L, Infante C, Parada SS, de Mayorga C. (2001) Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur J Orthod.* 23:153-167.
  20. Padma Kuman B, Retnakumari N. (2006) Loss of space and changes the dental arch after premature loss of the lower primary molar: a longitudinal study. *J Indian Soc Pedod Prev Dent.* 24:90-6.
  21. Nelson SJ, editor. *Wheeler's dental anatomy and occlusion.* 9th ed. 2009. Chapter 2.
  22. Leite-Cavalcanti A, Menezes SA, Granville-Garcia AF, Correia-Fontes LB. (2008) Prevalence of early loss of primary molars: study retrospective. *Acta Sci Health Sci.* 30:139-43.
  23. Cardoso L, Zembruiski C, Fernandes DS, Boff I, Pessin A. (2005) Evaluation of prevalence of precocious loss of deciduous molars. *Braz Res Pediatr Dent Integr Clin.* 5:17-22.
  24. Kelner N, Rodrigues MJ, Miranda K. (2005) Prevalence of early loss of deciduous molars in children attending the FOP/ UPE in 2002 and 2003. *Dent Clin Sci Recife.* 4:213-18.
  25. Al-Shahrani N, Al-Amri A, Hegazi F, Al-Rowis K, Al-Madani A, et al. (2015) The prevalence of premature loss of primary teeth and its impact on malocclusion in the Eastern Province of Saudi Arabia. *Acta Odontol Scand.* 73(7):544-9.
  26. *Orthodontics Australia.* "How genetics can affect your teeth". *Orthodontics Australia.* 2018-11-25.
  27. Corruccini RS, Potter RH. (1980) "Genetic analysis of occlusal variation in twins". *American J Orthodontics.* 78(2):140-54.
  28. Moimaz SA, Garbin AJ, Lima AM, Lolli LF, Saliba O, et al. (2014) "Longitudinal study of habits leading to malocclusion development in childhood". *BMC Oral Health.* 14(1):96.
  29. Graber TM. (1963) "The "Three m's": Muscles, Malformation and Malocclusion". *Am J Orthod.* 49(6):418-50.
  30. Liegeois F, Limme M. (1992) Space maintenance following the premature loss of temporary teeth. *Rev Belge Med Dent.* 47:9-22.
  31. Proffit WR, Fields HW, Moray LJ. (1998) Prevalence of malocclusion and orthodontic treatment need in the United States: estimates from the NHANES III survey. *Int J Adult Orthodon Orthognath Surg.* 13:97-106.
  32. Tausche E, Luck O, Harzer W. (2004) Prevalence of malocclusions in the early mixed dentition and orthodontic treatment need. *Eur J Orthodont.* 26:237-44.