

Effective Pulp Sensibility Tests Responses in Type 2 Diabetes Patients and Healthy Patients

Rahaf Marshad Almutairi, Deema Majid Alojayan, Noura Dshn Alajmi, Mzoon Mousa Alshehri* and Aseel Abdullah Ayidh

Department of Oral Biology, Riyadh Elm University, Riyadh, Saudi Arabia.

*Corresponding author: Mzoon Mousa Alshehri, Department of Oral Biology, Riyadh Elm University, Riyadh, Saudi Arabia.

Citation: Almutairi RM, Alojayan DM, Alajmi ND, Alshehri MM, Ayidh AA. (2022) Effective Pulp Sensibility Tests Responses in Type 2 Diabetes Patients and Healthy Patients. J Oral Med and Dent Res. 2(2):1-08.

Received: August 18, 2021 | **Published:** September 22, 2021

Copyright© 2021 genesis pub by Almutairi MR, et al. CC BY-NC-ND 4.0 DEED. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License., This allows others distribute, remix, tweak, and build upon the work, even commercially, as long as they credit the authors for the original creation.

Abstract

Diabetes mellitus (type II) is a metabolic disorder that affects the physiological functions of the body in a pernicious way. It has been shown that long term damage to vital organs including the oral cavity was brought on due to the elevation of blood glucose, which means that the structural components of the dental pulp may be directly affected as well as the influence on the sensory nerves too.

Keywords

Diabetes mellitus; Sensory nerves; Pulp sensibility test

Introduction

The dental pulp condition is assessed by stimulating the pulp through dental pulp testing, and that

makes it a useful and essential diagnostic tool. The pulp sensibility test includes thermal and electric tests, which determines pulp health from sensory response; A study was conducted on diabetes-induced rats where changes were noted in the structure and function of the blood vessels responsible for supplying the dental pulp [1].

Another study was conducted on diabetes mellitus (type II) patients, the results showed a significantly reduced response to the cold test when done on their upper premolars especially to those >45 years of age, the age of patients with diabetes might influence their response to the cold test [2].

In a study done by [3] pulp response to injury has been discussed; it being slower and more rapidly overwhelmed than in a non-diabetic tooth, and that studies have confirmed that the diabetic pulp has a lower blood flow. Estimation was done by World health organization (WHO) that around 7 million of the populations are diabetic even more concerning worldwide. Saudi Arabia ranks seventh in the world for the rate of diabetes [4].

Sensory disorders, taste problems, periodontal disease, xerostomia, salivary gland dysfunction, oral infections, and dental caries are oral problems that manifest in diabetic patients in high prevalence [5]. In addition, delayed mucosal wounds healing, as well as mucosal neuro-sensory disorders have been reported in patients with diabetes; there is a lack of awareness in regards to these complications worldwide [6]. Other than inducing dental pulp metabolic changes, diabetes mellitus has been shown to have the ability to influence both sensory and vascular structures [7]. Furthermore; little information exists regarding the responses of type II diabetes patients to the pulp sensibility tests (i.e., cold and electric) in Saudi Arabia. Therefore, the need of the study to be done will aid in improving the best intervention methods for diabetic patients.

The Hypothesis

The dental pulp response of type II diabetic patients differs than the pulp response of normal healthy patients.

Aims of the Study

To determine whether the pulp response of type II diabetic patients is slower than normal patients or is at the same rate.

Materials and Methods

This was an experimental study.

Study subject: diabetes patient (type 2) and normal healthy patient in the hospital RIYADH ELM UNIVERSITY, Riyadh, Kingdom of Saudi Arabia. Target sample size: Were around 100 male and female patients. Premolar teeth in 50 patients who had type 2 diabetes and Premolar teeth in 50 individuals with no medical condition were investigated. Electric and cold pulp sensibility tests were performed for premolar teeth and the electric pulp test results were recorded based on the pulp testers' grade that evoked a response. Power of the sample was calculated as well as the sample size. Duration of the

study: It will start at the beginning of February 2021 until beginning of Apr.

Methods: Electronic questionnaire was constructed, which was answered in the clinic after signing the informed consent. Random glucose test (blood sugar level around 200 milligrams or above per deciliter) and clinical trials pulp sensibility tests (cold and electric) were conducted. The collected data was subjected to statistical analysis using SPSS 22.

Inclusion:

- 51 diabetic (29 diabetic without any other medical condition and with medical condition) patients and 55 normal healthy patients were selected

Teeth should be:

- Premolars
- Sound; free of caries
- No recent history of trauma
- Vital

Exclusion:

- Patient with braces and pacemaker.
- Any other teeth except premolars
- Teeth with recent history of trauma
- Non vital teeth
- Carious teeth

The Questionnaire included:

Section 1: Demographic information

- Gender
- Age
- Medical condition.

Section 2: Disease information

- the duration of diabetes mellitus (type 2)
- the state of the patient diabetes mellitus (type 2)
- the result of random glucose test

Section 3: After this experimental

- patient response

Types of variables

- Nominal: gender, medical condition and state of the disease.
- Ordinal: Paine response.
- Ratio: age, the duration of disease and the result of random glucose test.

Based on the variables, the analysis types were done to find association and comparison

Results

A total of 106 cases were utilized in this study, which included 52 (49.1%) males and 54 (50.9%) females. Regarding their age, 8 (7.5%) belonged to 18-30 years, 36 (34%) belonged to 31-43 years, 38 (35.8%) to 43-55 years and 24 (22.6%) to 55 or above. Participants were also divided according to their medical condition, which showed that 29 (27.4%) had diabetes without any other medical condition, 22 (20.8%) had diabetes with another medical condition and 55 (51.9%) were healthy patients without any medical condition. Regarding their duration of diabetes, 31.4% had less than 5 years, 31.4% had 5-10 years and 37.3% had more than 10 years of duration. As far as their state was concerned, 86.3% had controlled and 13.7% had uncontrolled diabetes.

Pulp sensibility tests were conducted among all the participants and comparisons were made on the basis of gender, age, medical condition, duration of diabetes and state of diabetes on 4 different teeth of each study subject using Chi-square test. When compared on the basis of gender, statistically significant difference was observed at (p -value= .032) shown in table 1, however, no statistically significant differences between gender were found when conducted test on each teeth (p -value > .05). When compared on the basis of age, no statistically significant differences were found on each tooth (p -value > .05) as shown in table 2. When compared on the basis of duration of diabetes, statistically significant difference was obtained at (p -value = 0.039), while all other differences were not significant. Our main objective was to determine whether there is a difference in sensibility test results when conducted in patients with diabetes and without diabetes. It was noted from the findings that the patients having diabetes showed higher sensibility scores as compared to healthy patients. However, this difference was not statistically significant.

When compared the differences in sensibility test on the basis of duration of diabetes, statistically significant (p -value = 0.039) was found. It was noted that higher sensibility scores were recorded among patients with longest duration of diabetes as compared to shorter duration. Finally, the comparison between controlled and uncontrolled showed that there was no statistically significant found (p -value > .05) Figure 1 to Figure 3, Table 1 to table 3.

Gender Ratio






Figure 1: Gender Ratio.**Figure 2: Age distribution.****Figure 3: Medical condition.**

Item	Male	Female	P- Value
Gender	No Pain	No Pain	0.032
	12%	4%	
	Faint Pain	Faint Pain	
	2%	4%	
	Week Pain	Week Pain	
	27%	7%	
	Mild Pain	Mild Pain	
	23%	26%	
	Moderate Pain	Moderate Pain	
	16%	19%	
	Strong Pain	Strong Pain	
	6%	24%	
	Intense Pain	Intense Pain	
	12%	6%	
	Maximum Possible Pain	Maximum Possible Pain	
	2%	11%	

Table 1: Comparison on the basis of gender.

Item	18-30 Years	31-43 Years	44-55 Years	Above 55 Years	P- Value
Age	No Pain	No Pain	No Pain	No Pain	0.218
	25%	3%	8%	8%	
	Faint Pain	Faint Pain	Faint Pain	Faint Pain	
	13%	0%	3%	4%	
	Week Pain	Week Pain	Week Pain	Week Pain	
	0%	25%	13%	17%	
	Mild Pain	Mild Pain	Mild Pain	Mild Pain	
	13%	31%	32%	8%	
	Moderate Pain	Moderate Pain	Moderate Pain	Moderate Pain	
	0%	22%	18%	17%	
	Strong Pain	Strong Pain	Strong Pain	Strong Pain	
	38%	11%	8%	25%	
	Intense Pain	Intense Pain	Intense Pain	Intense Pain	
	0%	3%	5%	4%	
	Maximum Possible Pain	Maximum Possible Pain	Maximum Possible Pain	Maximum Possible Pain	
	13%	6%	13%	17%	

Table 2: Comparison on the basis of age.

Discussion

This study aimed to determine the effect of diabetes on the results of sensibility test, which has resulted in finding that overall there was no statistically significant difference among diabetic and healthy

patients. Moreover, no association of age and the effect of sensibility test on the tooth were observed in our study. A similar study conducted by Kermani MT, et al. [2] revealed a statistically significant association of age with the pulp response to sensibility tests. Moreover, they also found statistically significant difference between diabetic and healthy patients, which reported that the diabetic patients aged more than 45 years showed the least number of pulpal responses as compared to younger age groups and healthy patients. These findings were found to be different from what we observed among our study participants.

Another similar study done by Moderasi, et al. [8] revealed that there was difference between diabetic group and healthy patients group. Still, after performing the test, the association was statistically insignificant. The achieved difference in results was reliable considering the size of sample. Additionally, findings of this study disclose the correlation between age and electrical stimulation threshold of teeth. A small difference between age groups was detected in regard to electrical stimulation threshold of teeth. This indicates that increased age leads to decreased sensitivity of teeth due to decreased size of pulp chamber. These findings are dissimilar to what we found, as both age and medical conditions were not statistically significant associated with the sensibility scores.

One more investigation done by Barczak, et al. [9] reported that the threshold was significantly lesser in younger patients. The threshold of pulp sensitivity of the remaining groups of teeth was similar in both age groups. The correlation between sensibility of the pulp in the Caucasian population and gender was not confirmed. Age related findings are not similar to our study but the gender association is similar to what we found. However, there is a major disparity between Barczak, et al. [9] and Moderasi, et al. [8] when compared the age groups as former study reported lower sensibility scores among younger age group and later among the older age groups. This is interesting to know as the later study also involved diabetic patients in their investigation, which may play an important role in determining this causal relationship. However, more studies need to be done in order to find a stronger association (if any) between diabetes and sensibility test scores [10,11].

References

1. Fenn S, Narayanan M, Jacob M. (2019) Insidious Role of Diabetes Mellitus on Nerves and Dental Pulp. *J Clinic Diag Res.* 13(3):5-7.
2. Kermani MT, Sanjari M, Nakhaei N, Parirokh M, Abbottn P, et al. (2020) Comparison of Pulp Sensibility Tests Responses in Type 2 Diabetes Patients and Healthy Individuals. *J Endod.* 46(3):364-9.
3. Moraru AI, Gheorghijă LM, Dascălu IT, Bătăiosu M, Manolea HO, et al. (2017) Histological and immunohistochemical study on the dental pulp of patients with diabetes mellitus. *Rom J Morphol Embryol.* 58(2): 493-9.
4. Al Dawish Abdulaziz M, Alwin Robert A, Braham R, Abdallah Al Hayek A, Al Saeed A, et al. (2016) Diabetes mellitus in Saudi Arabia: a review of the recent literature *Curr Diabetes Rev.* 12(4):359-8.
5. Nazir MA, AlGhamdi L, AlKadi M, AlBeajan N, AlRashoudi L, et al. (2018) The burden of diabetes, its oral complications and their prevention and management. *Open Access Maced J Med Sci.* 6(8):1545-53.
6. Al-Maskari AY, Al-Maskari MY, Al-Sudairy S. (2011) Oral manifestations and complications of diabetes mellitus: a review. *Sultan Qaboos Univ Med J.* 11(2):179-6.

7. Ciesielski M, Mochnacki B, Siedlecki J. (2016) Simulations of thermal processes in tooth proceeding during cold pulp vitality testing. *Acta Bioeng Biomec.* 18(3):33-1.
8. Modaresi J, Afkhami-Ardekani M, Mokhtari F, Salmani-Qahyazi A, Amirzade-Iranaq MH. (2017) The comparison of tooth pulp electrical threshold in patients with type II uncontrolled diabetes and healthy subject. *Iranian Journal of Diabetes and Obesity.* 9(1):20-4.
9. Barczak K, Palczewska-Komsa M, Wilk A, Nowicka A, Buczkowska-Radlińska J, et al. (2020) Pulp sensibility to electric stimuli in the Caucasian population. *Aust Endod J.* 46(1):26-2.
10. Rowe HA, Pitt Ford TR. (1990) "The assessment of pulpal vitality,". *Int Endod j.* 23(2):77-3.
11. Catanzaro O, Dziubecki D, Lauria LC, Ceron CM, Rodriguez RR. (2006) Diabetes and its effects on dental pulp. *J Oral Sci.* 48(4):195-9.