

## Reducing Osseointegration Time using UV Machine Implants Treatment: Clinical Report

Angelo Cardarelli\*

Oral Surgeon, Adjunct Professor UCAM University of Spain, Gargash Hospital Dubai Clinica, Cardarelli, Italy, Dona Medical Clinic, Dubai

\***Corresponding author:** Oral Surgeon, Adjunct Professor UCAM University of Spain, Gargash Hospital Dubai Clinica, Cardarelli, Italy, Dona Medical Clinic, Dubai.

**Citation:** Cardarelli A. (2023) Reducing Osseointegration Time Using UV Machine Implants Treatment: Clinical Report. J Oral Med and Dent Res. 4(2):1-10.

**Copyright**©2023 by Cardarelli A. All rights reserved. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Received:** July 03, 2023 | **Published:** July 21, 2023

### Abstract

This study's goal is to determine how well UV therapy works for placing dental implants. Panoramic X-ray, clinical image, CBCT were used to assess the effects of UV irradiation on reversing the biological aging process of implants. BIC is predominantly influenced by implant surface characteristics. Among the most popular methods of modifying an implant surface is the roughening of the surface using the sandblasted, large-grit, acid-etched (SLA) technique.

### Keywords

Ultraviolet; Dental implant; Osseointegration; Bone-implant interface.

### Introduction

The most important factor for osseointegration of implants is the correct interface between macro- and micro-design, which varies depending on the classification of implants into three types based on external bone level, internal tissue level, and internal bone level. The implant surface has an

approximately 50-nm-thick TiO<sub>2</sub> oxide film. With time, hydrocarbons in the air are adsorbed on to the oxide film, leading to a biologic aging process that prevents osseointegration. [1-2-3-4] The surface of implant begins to change from hydrophilic to hydrophobic after the manufacturing process. Efforts have been made to reverse this biological aging process and to induce the original hydrophilic state through UV irradiation. In line with this trend, in the present report, I will share my experience with UV machine (AB DENTAL) [5-10].

## Materials and Methods

UV photo functionalization was performed by illuminating implants with UV light for 15 seconds using a specialized instrument (AB DENTAL UV ACTIVATOR ISRAEL). Specifications of the device were as follows: input voltage of AC 100 to 240 V  $\pm$  10%, input current of 2.2 A in maximum. Multiple UV lamps were utilized to create the UV light as a mixed spectrum at wavelengths of 172 nm. Inside the device, the lamps were arranged to illuminate an implant in all directions homogeneously. We have used Ab dental implants that have had their surfaces roughened with calcium phosphate particles before undergoing multiple thermal acid etching procedures to determine the microstructure and surface roughness of the implants. Then, gamma irradiation is used to sterilize all implants.



**Figure 1:** Implants surface after UV treatment increasing the Hydrophilicity (AB DENTAL).



**Figure 2:** UV machine (AB DENTAL).

## First Case: Immediate Implant in Molar Area

Female patient 45 years old.



**Figure 3:** Immediate implant placement in mandibular molar area; Extraction and implant (I55 4,2 x11,5 Ab dental) and dentin graft.



**Figure 4:** Healing after 2 months and temporary restoration.



**Figure 5:** X-ray after 2 months.

## Second Case: Immediate Implants and Crestal Sinus Lift

Male patient 72 years old.

Extraction and immediate implants (AB dental I5) 3,75 x13 and 4,2 x 11,5 and crestal sinus lift with osseodensification using dentin GRAFT IVORY GRAFT ISRAEL.



**Figure 5:** Extraction and immediate implants.



**Figure 6:** X-ray Initial.



**Figure 7:** X-ray after implants placement.



**Figure 8:** Healing after 3 months and temporary restoration.




Figure 9: CBCT after 3 months.

### Third Case: Immediate Implants and immediate restoration

Female patient 79 years old extraction and immediate implants (AB dental) dentin graft (IVORY GRAFT).



Figure 10: Initial X-ray.





Figure 11: Extraction and immediate implants.




Figure 12: Immediate restoration.



**Figure 13:** X-ray after implants placement.



**Figure 14:** Healing after 3 months.



**Figure 15:** CBCT after 3 months.

#### Fourth Case: Immediate Implant in Esthetic Area with Immediate Restoration

Male Patient 50 Years old extraction immediate implant Dentin Graft (IVORY GRAFT) and immediate restoration.



**Figure 16:** Extraction and immediate implant with Dentin Graft.



**Figure 17:** Extraction immediate implant Dentin Graft.



**Figure 18:** Immediate temporary restoration



**Figure 19:** Immediate temporary restoration.



**Figure 20:** Final restoration after 3 months.



**Figure 21:** X-ray after 3 months



**Figure 22:** CBCT 3 months later



## Discussion

Photo functionalization are due to improving hydrophilicity and eliminating hydrocarbon contamination on the implant surface in addition to antibacterial effects, UV activation would increase the adsorption of plasma proteins of human body and improve osteogenic cell attachment, spreading, and proliferation. There is an important bacterial colonization on implants only 30 min after implant insertion which may be prevented by UV-photofunctionalization restricting the growth of oral bacteria and biofilm. Peri-implant-diseases-associated biofilms would affect the long-term outcome of dental implants. The microbial composition between periodontitis and peri-implantitis are similar; however, dental implants are more susceptible to oral infections than natural teeth [11-14].

## Conclusion

Many methods have been used to promote osseointegration for successful implant therapy, including photofunctionalization. UV photofunctionalization can change the surface wettability and eliminate the hydrocarbons that are generated by aging on the implant surface [15-17]. According our clinical experience using this machine we have a significative reduction in terms of osseointegration time also in case of immediate restoration and post extractive procedures. However, further studies must be conducted in the future.

## References

1. Funato A, Ogawa T. (2013) Photofunctionalized dental implants: A case series in compromised bone. *Int J Oral Maxillofac Implant.* 28(6):1589-601.
2. Funato A, Yamada M, Ogawa T. (2013) Success rate, healing time, and implant stability of photofunctionalized dental implants. *Int J Oral Maxillofac Implant.* 28(5):1261-71.
3. Suzuki S, Kobayashi H, Ogawa T. (2013) Implant stability change and osseointegration speed of immediately loaded photofunctionalized implants. *Implant Dent.* 22(5):481-90.
4. Kitajima H, Ogawa T. (2016) The Use of Photofunctionalized Implants for Low or Extremely Low Primary Stability Cases. *Int J Oral Maxillofac Implant.* 31(2):439-47.
5. Hirota M, Ozawa T, Iwai T, Ogawa T, Tohnai I. (2016) Implant Stability Development of Photofunctionalized Implants Placed in Regular and Complex Cases: A Case-Control Study. *Int J Oral Maxillofac Implant.* 31(3):676-86.
6. Hirota M, Ozawa T, Iwai T, Ogawa T, Tohnai I. (2018) Effect of Photofunctionalization on Early Implant Failure. *Int J Oral Maxillofac Implants.* 33(5):1098-1102.
7. Hirota M, Ozawa T, Iwai T, Mitsudo K, Ogawa T. (2020) UV-Mediated Photofunctionalization of Dental Implant: A Seven-Year Results of a Prospective Study. *J Clin Med.* 9(9):2733.
8. Puisys A, Schlee M, Linkevicius T, Petrakakis P, Tjaden A. (2020) Photo-activated implants: A triple-blinded, split-mouth, randomized controlled clinical trial on the resistance to removal torque at various healing intervals. *Clin Oral Investig.* 24(5):1789-1799.
9. Choi B, Lee YC, Oh KC, Lee JH. (2021) Effects of photofunctionalization on early osseointegration of titanium dental implants in the maxillary posterior region: A randomized double-blinded clinical trial. *Int J Implant Dent.* 7(1):37.
10. Shah SA, Singh BP, Rao J, Kumar L, Singh M, et al. (2021) Biological and esthetic outcome of immediate dental implant with the adjunct pretreatment of immediate implants with platelet-rich plasma or photofunctionalization: A randomized controlled trial. *J Indian Prosthodont Soc.* 21(4):348-355.

11. Tominaga H, Matsuyama K, Morimoto Y, Yamamoto T, Komiya S, et al. (2019) The effect of ultraviolet photofunctionalization of titanium instrumentation in lumbar fusion: A non-randomized controlled trial. *BMC Musculoskelet Disord.* 20(1):292.
12. Pesce P, Menini M, Santori G, Giovanni E, Bagnasco F, et al. Photo and Plasma Activation of Dental Implant Titanium Surfaces. A Systematic Review with Meta-Analysis of Pre-Clinical Studies. *J Clin Med.* 9(9):2817.
13. Dini C, Nagay BE, Magno MB, Maia LC, Barao VAR. (2020) Photofunctionalization as a suitable approach to improve the osseointegration of implants in animal models-A systematic review and meta-analysis. *Clin Oral Implant Res.* 31(9):785-802.
14. Yeo IL. (2019) Modifications of Dental Implant Surfaces at the Micro- and Nano-Level for Enhanced Osseointegration. *Materials (Basel).* 13(1):89.
15. Roy M, Pompella A, Kubacki J, Szade J, Roy RA, et al. (2016) Photofunctionalization of Titanium: An Alternative Explanation of Its Chemical Physical Mechanism. *PLoS One.* 11(6):0157481.
16. Roy M, Pompella A, Kubacki J, Piosik A, Psiuk B, et al. (2017) Photofunctionalization of dental zirconia oxide: Surface modification to improve bio-integration preserving crystal stability. *Colloids Surf B Biointerfaces.* 156:194-202.
17. Arroyo-Lamas N, Ugalde U, Arteagoitia I. (2020) Decontamination of Ti Oxide Surfaces by Using Ultraviolet Light: Hg-Vapor vs. LED-Based Irradiation. *Antibiotics (Basel).* 9(11):724.