

Curious Case of an Anterior Neck Swelling

Shaikh Mohd Shaad^{1*}, Mehta Keshavi¹, Gajghate Nayan¹ and Sanjay Chatterjee²

¹Post Graduate Student, Department of General Surgery, Bombay Hospital & Institute of Medical Sciences, Mumbai, India

²A Assistant Professor, Maharashtra University of Health Sciences, Nashik Postgraduate Teacher, Bombay Hospital Institute of Medical Sciences, Bombay Hospital Avenue, New Marine Lines, Mumbai, India

* **Corresponding author:** Chatterjee S, Post Graduate Student, Department of General Surgery, Bombay Hospital & Institute of Medical Sciences, Mumbai, India.

Citation: Shaad SM, Keshavi M, Nayan G, Chatterjee S. (2023) Curious Case Of An Anterior Neck Swelling. Genesis J Surg Med. 2(2):1-6.

Received: November 06, 2023 | **Published:** November 22, 2023

Copyright © 2023 genesis pub by Shaad SM, et al. CC BY-NC-ND 4.0 DEED. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License. This allows others distribute, remix, tweak, and build upon the work, even commercially, as long as they credit the authors for the original creation.

Abstract

A Thyroglossal cyst is amongst the most commonly encountered anterior painless neck swelling. Its usual presentation is seen in childhood and rarely it can present in adulthood. The management for Thyroglossal cyst is, the Sistrunk's Operation. The etiology is explained by embryonic development of the Thyroglossal duct and its deficiency to obliterate after birth Here we present a case of a 45-year-old gentleman who presented with a painless midline neck swelling for 6 years. Onset of the swelling was insidious and gradually increased in size over the years. Patient did not give any history suggestive of an infection or thyroid gland dysfunction. USG of the neck revealed a fluid filled cystic lesion below the hyoid bone. MRI of the Neck revealed a 3.5 x 3 x 3.5 cm well-defined lesion in the anterior neck suggesting a Thyroglossal Cyst. Patient was worked up for a Sistrunk's procedure. Histopathology revealed a well-differentiated Papillary Thyroid Carcinoma.

Keyword

Thyroglossal cyst; Thyroglossal duct; Sistrunk's procedure; Papillary thyroid carcinoma; Malignancy

Introduction

Thyroglossal duct cyst (TDC) is one of the most common midline swellings of the neck which arises from an unobliterated thyroglossal duct which is present in the embryonic period [3]. It represents the embryonic pathway of descent of the thyroid gland. As it is a congenital disease it presents most commonly in the childhood period and seldom presents in adulthood. Hence, most of the data available on this subject have a focus on management of this condition in the childhood period.

Sistrunk's Procedure is the procedure of choice for this disease and is accepted globally as the standard of care [4]. The operation includes the excision of the cyst with excision of the central part of the hyoid bone and coring of the tract till the foramen cecum. This is driven by the knowledge of the embryonic development of the Thyroglossal duct and its intimate relation with the Hyoid bone.

Data available reveals a satisfactory cure rate with Sistrunk's Operation for Thyroglossal cyst occurring in childhood and very few cases are reported in adult age group and hence less data is available for comparison [6].

Case report

Here we present a case of a 45-year-old gentleman with no comorbidities, who presented with a painless midline neck swelling since 6 years, the swelling was first noticed by the patient 6 years ago when it was the size of a pea gradually it progressed to the current size of a lemon. Patient did not give any history of Fever, Pus discharge from swelling. No history suggestive of thyroid dysfunction or difficulty in swallowing/breathing/change of voice. Examination revealed a 4 cm x 4 cm midline neck swelling, moving with deglutition and protrusion of tongue, smooth surface with rounded edges, no tenderness or rise of local temperature. Thyroid gland not palpable. No cervical lymph nodes palpated.



Figure 1: Movement with tongue protrusion.



Figure 2: Midline neck swelling.

MRI Neck revealed well defined T1 hyper intense swelling in the infra hyoid region with multiple blooming spots suggestive of thyroglossal cyst with calcifications.




Figure 3: MRI neck in sagittal view.




Figure 4: Antero-posterior view showing close relation the hyoid bone.

CECT of Neck revealed a well-defined peripherally enhancing cystic structure in midline upper neck infra hyoid in location measuring 3.6 cm x 2.9 cm x 3.5 cm with foci of calcification.




Figure 5: Image showing calcifications.

Patient was posted for Sistrunk's procedure. Patient, under general anaesthesia, was taken in the supine position with neck extended and head supported with the help of a ring. A transverse incision was taken over the swelling and deepened till the investing layer of deep cervical fascia. Strap muscles were retracted laterally and swelling was freed from all adhesions. Hyoid bone was located superiorly to the swelling, off which, the muscle attachments from the central part were released. The hyoid bone was cut in the central part. The tract was then seen extending superiorly into floor of the mouth and cored upto the mylohyoid muscle. Wound was closed in layers after achieving homeostasis over a closed vacuum drain. Patient tolerated the procedure well and the specimen was sent for a Histopathological Exam. Post operative period was uneventful and the patient was discharged of POD-2 with drain in-situ.

Histopathological examination revealed a well-differentiated Papillary Thyroid Carcinoma involving the capsule with classical orphan Annie nuclei.




Figure 6: Papillary fronds with orphan- annie nuclei noted.

Discussion

The rare occurrence of thyroglossal cyst in adults is the reason for the paucity of data in the adult population. The majority of patients are males. Reports in literature, however, do not come to a consensus with some reports suggesting equal distribution, whilst some suggest male or female preponderance.

Painless anterior midline neck swelling is the most common presentation of Thyroglossal Cyst in adult patients. The presence of other symptoms such as pain, odynophagia, dysphagia, and dyspnoea often indicates the presence of complications such as abscess formation. Location of thyroglossal cyst with respect to hyoid bone can be variable, while midline position is the dominant location in both children and adults, lateral deviation has been noted in adult presentation [3]. With respect to the hyoid bone, cysts can be above, over, or below the hyoid, most commonly they are infra-hyoid. The surgical management of thyroglossal cyst has changed with time. Earlier Incision & Drainage or simple cyst excision were presented with unacceptably high levels of recurrence, Schlange in 1893, suggested excision of the cyst and mid-portion of the hyoid bone and leaving behind the proximal tract – a technique which resulted in recurrence rates of 30 %. In 1920 Walter Ellis Sistrunk reported the classical Sistrunk procedure which significantly improved postoperative outcomes and has since remained the gold standard to date [2]. Recent advancements such as robot-assisted, endoscope-assisted transoral, axillary & retro-auricular approaches have been attempted for thyroglossal cyst in adults. These procedures are cosmetically superior to the sistrunks operation but data on efficacy being the same are inadequate. Recurrence is the most important post-op outcome following the Sistrunk procedure, with a recurrence noted in 3%–6% of cases. This is often attributed to technical shortcomings viz. incomplete excision of the duct or the presence of ramification of ducts, which remain unrecognized at the time of surgery. Most recurrences occur in the first 6 months [6]. Another important possible complication is damage to the 12th Cranial nerve. Meticulous dissection of the central part of hyoid bone and preservation of the Superior Horn of Hyoid bone can help significantly in preventing injuries to the Hypoglossal nerve.

Thyroglossal Cyst present with a <1% chance of harboring a malignant focus in adult population [1]. Most commonly on histopathology we find papillary thyroid carcinoma (90 %) (5). Many treatment options such as Total Thyroidectomy with Bilateral neck dissection, radioactive iodine, thyroid suppression have been suggested. Sistrunks operation is usually considered sufficient and need for further intervention is decided on the basis on the following findings a.) Thyroid nodule (cold) picked up on thyroid scan b.) cervical lymph nodes detected clinically or on imaging c.) prior history of irradiation to the neck. Calcification is the hallmark of papillary carcinoma in a thyroglossal duct cyst [5]. Pre-operative role of FNAC remains uncertain but should be recommended to pick up lesions early and to plan a definitive treatment [6]. The Sistrunk operation alone is sufficient for squamous carcinoma, but total thyroidectomy is recommended for differentiated thyroid carcinoma.

After lingual thyroid, the Thyroglossal cyst in the second most-common site for ectopic thyroid tissue.

Incidence of which is approximately 60%. In most of the cases the thyroid gland is normal and the patients are often euthyroid or hypothyroid at presentation. Hence, pre-operative evaluation with a thyroid function test and thyroid scan should be undertaken, prior to its excision, so as to confirm whether the tissue in the cyst is not the only functioning thyroid tissue in the body.

References

1. Dedivitis RA, Guimarães AV (2000). Papillary thyroid carcinoma in thyroglossal duct cyst. *Int Surg.* 85(3):198–201.
2. Allard RH. (1982) The thyroglossal cyst. *Head Neck Surg.* 5(2): 134–46.
3. Weiss SD, Orlich CC. (1991) Primary papillary carcinoma of a thyroglossal duct cyst: report of a case and literature review. *Br J Surg.* 78(1):87–9.
4. Taori K, Rohatgi S, Mahore DM, Dubey J, Saini T. (2005) “Papillary carcinoma in a thyroglossal duct cyst” - a case report and review of literature. *Indian J Radiol Imaging.* 15(4):531–33.
5. Vera-Sempere F, Tur J, Jaén J, Perolada JM, Morera C. (1998) Papillary thyroid carcinoma arising in the wall of a thyroglossal duct cyst. *Acta Otorhinolaryngol Belg.* 52(1):49–54.
6. Yang YJ, Haghiri S, Wanamaker JR, Powers CN. (2000) Diagnosis of papillary carcinoma in a thyroglossal duct cyst by fine-needle aspiration biopsy. *Arch Pathol Lab Med.* 124(1):139–42