

The Thalamic Oscillatory Integration Theory: A Unified Framework for Spatial Orientation and Balance TOIT: A Conjectural Extension of Default Space Theory to Vestibular Function and Postural Control

Ravinder Jerath¹ and Varsha Malani^{2*}

¹Conceptualization, Data curation, Project administration, Visualization, Writing – original draft

²Investigation, Methodology, Software, Visualization, Writing – review & editing

* **Corresponding author:** Varsha Malani, Investigation, Methodology, Software, Visualization, Writing – review & editing

Citation: Jerath R, Malani V The Thalamic Oscillatory Integration Theory: A Unified Framework for Spatial Orientation and Balance TOIT: A Conjectural Extension of Default Space Theory to Vestibular Function and Postural Control. J Neurol Sci Res. 6(2):1-16.

Received: September 23, 2026 | **Published:** October 05, 2026

Copyright© 2026 Genesis Pub by Jerath R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0). This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are properly credited.

Abstract

The Default Space Theory (DST) of consciousness holds that conscious experience arises from synchronized physiological and neural oscillations that construct an internal three-dimensional “default space” approximating the body’s position in the external world. Here we propose the Thalamic Oscillatory Integration Theory (TOIT), a conjectural extension of DST to spatial orientation and postural balance. We propose that fast gamma-band oscillations (~30–100 Hz) bind visual, auditory, somatosensory, and motor input into a coherent internal spatial map, while slower thalamic theta (4–8 Hz) and alpha (8–12 Hz) rhythms provide temporal orientation — gating sensory input and permitting rapid postural correction. We synthesize anatomical, histological, and electrophysiological findings from peripheral receptors (retina, cochlea, vestibular apparatus) and central structures (cerebellum, thalamus, cortex) into a single oscillatory account, and contrast it with the conventional model, in which the cerebellum performs error-correction on vestibular and proprioceptive feedback.

Under TOIT, thalamic integrative gating and cross-modal gamma synchrony play the larger role — a claim we argue is consistent with clinical syndromes such as vestibular migraine and sensory ataxia, where balance fails despite intact

cerebellar circuitry. We outline testable predictions (e.g., increased theta-gamma coupling during active navigation) and potential interventions (e.g., 40 Hz or theta-frequency neural entrainment) that would help distinguish TOIT from the standard model rather than merely being compatible with it.

Keywords

Vestibular system; Spatial representation; Brain oscillations; Consciousness; Spatial orientation; Dynamic connectivity.

Introduction

Maintaining orientation and upright posture is something people do constantly without conscious effort — while standing still, walking, climbing stairs, or dancing to an unpredictable rhythm. That ease is itself notable. A process this automatic, this continuously error-corrected, and this robust to unexpected perturbation is usually assumed to require dedicated, specialized hardware; a robot attempting the same feat needs an explicit sensor-fusion algorithm running many times per second. Yet subjectively, human balance requires no attention at all until something goes wrong — a slick patch of floor, a missed stair, a rocking boat deck — at which point correction is not merely fast but appears to occur before the disturbance reaches awareness.

The Default Space Theory (DST), developed by Jerath and colleagues since 2015, proposes that this ease reflects a deeper property of consciousness itself: the brain and body continuously construct a three-dimensional internal simulation (“default space”) of the surrounding world through synchronized oscillations across sensory and motor systems [1]. In this view, perception is not a passive readout of the world but an actively maintained internal model, continuously updated and continuously compared against incoming sensory evidence. The thalamus is treated as a central integrating hub in this process — fast oscillatory activity converging from the eyes, ears, and skin is proposed to be bound there into a continuously updated three-dimensional reference frame [2]. The theory has since been extended with phenomenological analysis of subjective spatial experience [2] a discussion of some of consciousness’s enduring puzzles through this lens [3], a broader proposal about the hierarchical organization of oscillatory assemblies across scales [4], and a proposed account of how reflexive, sub-cortical mechanisms bind into conscious experience [5].

This paper extends DST into a framework we term the Thalamic Oscillatory Integration Theory (TOIT), applied specifically to spatial orientation and balance — domains conventionally attributed almost entirely to the vestibular system and cerebellum, with the thalamus cast in a supporting, largely relay-like role. We propose instead that gamma-band synchrony across visual, auditory, and somatosensory cortex constructs an internal spatial map that substitutes, moment to moment, for direct perception of the external world, and that slower thalamic rhythms — theta (~4–8 Hz) and alpha (~8–12 Hz) — provide the timing and gating needed to keep that map aligned with incoming sensory data. Hippocampal theta oscillations, already well established in encoding spatial location via place cells and grid cells [6,7], are proposed here to serve a parallel role in fusing visual distance information with vestibular directional cues — an extension by analogy from a well-documented mechanism in one domain (episodic spatial memory)

to a different but related domain (moment-to-moment postural orientation) that has not, to our knowledge, been directly tested.

This account departs from standard textbook models, which locate balance control primarily in a cerebellar reflex arc: vestibular and proprioceptive input is relayed through the brainstem to the cerebellum, which computes and corrects motor error, with the thalamus serving mainly as a relay station [8]. We argue this relay-only view understates the thalamus's contribution. Evidence for direct thalamic integration of vestibular signals — including thalamic neurons that respond to head orientation independent of downstream cortical processing — suggests the thalamus does more than pass information along unmodified [11]. Clinically, this distinction is not merely semantic: patients with intact cerebellar function can still lose the ability to orient themselves after thalamic damage [14], a finding that is difficult to explain under a purely cerebellum-centered model but is at least consistent with, and arguably predicted by, a thalamus-centered one.

The chapters that follow lay out: (1) the standard model and its limitations; (2) the TOIT framework as applied to orientation and balance; (3) the relevant anatomy; (4) known oscillatory phenomena across each sensory system; (5) the proposed thalamic dual-oscillator mechanism; (6) cross-frequency coupling as the proposed integrating mechanism; (7) how this account maps onto clinical vestibular and balance disorders; (8) a concrete methodological roadmap for testing the theory's most specific claims; and (9) the limitations of the present account, stated plainly rather than left implicit.

Prevailing Model: Vestibular–Cerebellar Balance Control

Standard accounts treat orientation and balance as a hierarchical sensorimotor reflex loop, and it is worth being fair to how much this model explains before turning to its gaps. The vestibular apparatus — the semicircular canals and otolith organs — detects head movement and gravitational orientation and signals the brainstem vestibular nuclei, triggering compensatory reflexes on a timescale of tens of milliseconds. Proprioceptive input from muscles and joints provides parallel information about limb and trunk position, arriving via a separate but converging pathway. The cerebellum integrates both streams, calibrating output to correct posture and gait by modulating muscle tone and stretch-reflex gain [8,9]. The vestibulocerebellum in particular (the flocculonodular lobe) stabilizes eye movements and axial posture; damage to the cerebellar vermis reliably produces truncal ataxia and a wide-based gait [10], which is among the clearest structure–function relationships in all of motor neuroscience.

In this model, detected sway is corrected through vestibulo-spinal and stretch reflexes, with the cerebellum minimizing error iteratively — essentially performing a continuous forward-model comparison between predicted and actual sensory consequences of movement. The thalamus is treated as a relay for conscious awareness of tilt or motion, not as part of the feedback loop itself [11], and multisensory (visual–vestibular) integration is generally attributed to cortical regions such as the parietal “vestibular cortex” or the temporo-parietal junction, which are thought to construct a subjective sense of self-motion and orientation downstream of the reflexive correction already performed by the brainstem and cerebellum.

This relay-only view, however, is increasingly hard to square with newer anatomical and physiological

findings. Vestibular nuclei project directly onto thalamic nuclei, which integrate vestibular, visual, and somatosensory input before signals ever reach cortex [12] — a wiring pattern that would be superfluous if the thalamus's only job were passive relay. In rodents, neurons in the anterior and laterodorsal thalamus encode head direction from combined vestibular and visual input, functioning as something closer to an internal compass than a passive relay station [13,12]. Even the simple act of standing requires reconciling conflicting sensory streams — on a moving ship, for instance, vision reports stillness while the vestibular system reports motion — and this reconciliation process appears to recruit cortical oscillatory networks alongside the cerebellum, particularly gamma-band synchrony thought to bind multisensory information together [13].

Clinical cases sharpen the point further. Thalamic stroke can produce “thalamic astasia” — an inability to maintain upright posture with no vestibular or motor deficit whatsoever [14]. Vestibular migraine patients show abnormal thalamic activation in response to vestibular stimulation despite entirely normal peripheral vestibular function [15]. Both observations are difficult to explain if the thalamus is merely a relay station passing signals through unmodified. We take this as motivation for a model — TOIT — in which the thalamus plays an active, integrative role in constructing and maintaining spatial orientation, rather than a passive one in merely reporting it to consciousness after the fact.

The Thalamic Oscillatory Integration Theory Framework

TOIT holds that consciousness and spatial self-location emerge from interconnected oscillations across brain and body, producing a “default space” that structures both perception and action [1]. Gamma oscillations (~30–100 Hz) are well established as a mechanism for binding features within a single sensory modality — for example, binding color, shape, and motion into a unified visual object. TOIT proposes that the same binding mechanism operates across modalities as well: for instance, aligning visual space with head orientation via visual–vestibular gamma synchrony [4]. This is proposed to produce an internally maintained three-dimensional space that persists briefly even without continuous external input [2] — a property TOIT also invokes, more speculatively, to explain dreaming, treating dream imagery as internally generated content filling the same default space in the absence of external sensory transduction.

Within this framework, TOIT assigns the thalamus an organizing role via two distinct rhythms operating in parallel. Theta (~4–8 Hz) is linked to anterior thalamic and hippocampal circuits, and is proposed to provide a recurring “timestamp” that aligns incoming vestibular and proprioceptive signals with the current internal spatial model — in effect, a periodic checkpoint at which the brain asks whether its running simulation still matches the body's actual position. Alpha (~8–12 Hz), linked to reticular and sensory relay nuclei, is proposed to gate which sensory information reaches conscious awareness at all, filtering out noise — minor postural sway, ordinary visual jitter — so that only genuine perturbations register and demand a correction.

Under this framework, standing on one foot involves continuous micro-corrections that occur without conscious involvement because the thalamus is constantly comparing incoming signals against the internal model and correcting deviations before they reach awareness at all — which would explain why balance feels effortless right up until the moment a correction is large enough, or sudden enough, to break through the alpha-mediated gate and become a conscious experience of “almost falling.”

TOIT further proposes that this oscillatory activity is not confined to canonical “processing” regions of cortex but is distributed across cellular and tissue-level electrical signaling more broadly [1,4]. This is, by the authors’ own framing, the most speculative part of the theory and the hardest to test directly, since it makes a claim about bioelectric activity at a scale well below what current human-subject recording methods can resolve non-invasively. We flag it here explicitly as a hypothesis awaiting technique development, not as an established mechanism, and we return to this limitation in Section 10.

In summary, TOIT proposes a layered structure: gamma-band synchrony constructs moment-to-moment spatial detail; thalamic theta and alpha rhythms provide timing and gating for that detail; and the resulting default space remains stable until a genuine perturbation — a loss of balance, a sudden movement, an unexpected visual shift — forces a correction that is large enough to surface into conscious awareness.

Anatomical and Histological Foundations

Visual system

Retinal photoreceptors transduce light into electrochemical signals that travel via the optic nerve to the thalamic lateral geniculate nucleus (LGN) and then onward to primary visual cortex (V1), along parallel parvocellular and magnocellular channels specialized respectively for fine detail/color and motion/contrast. Notably, only about 10% of synapses onto LGN relay cells originate from the retina itself; the remaining ~90% come from cortical feedback and local interneurons [16]. This anatomical asymmetry is one of the stronger pieces of general evidence that the LGN — and by extension the thalamus more broadly — is not a passive relay but actively shapes signals in light of ongoing cortical context, a structural fact TOIT leans on heavily in extending the same logic to vestibular and somatosensory thalamic nuclei.

Auditory system

Hair cells in the organ of Corti transduce sound vibration into action potentials that travel via the cochlear nuclei to the medial geniculate nucleus (MGN) and then to primary auditory cortex (A1), organized tonotopically at every stage of the pathway. Outer hair cells also produce active mechanical oscillations of their own (otoacoustic emissions), and cortically, gamma-band activity in A1 has been linked to auditory feature binding in a manner directly analogous to visual gamma binding [17].

Vestibular system

The semicircular canals (rotation) and otolith organs (linear acceleration, tilt) transduce head movement into signals carried by cranial nerve VIII to the brainstem vestibular nuclei. From there, pathways ascend to the cerebellum via the flocculonodular lobe, descend via vestibulospinal tracts to spinal motor neurons, and — critically for TOIT — ascend through the thalamus, primarily via the ventral posterolateral (VPL) and posterior nuclei, which also process somatosensory input and are therefore a natural site for vestibular–somatosensory convergence [18]. Head-direction cells in thalamic and limbic structures fire according to head orientation relative to gravity [19], and thalamic projections to parieto-insular vestibular cortex (PIVC/OP2) integrate vestibular, proprioceptive, and visual signals for real-time spatial localization, providing an anatomical substrate for exactly the kind of cross-modal convergence TOIT’s gamma-synchrony proposal requires.

Somatosensory system

Proprioceptive and cutaneous signals travel via the dorsal column–medial lemniscal pathway to thalamic VPL/VPM nuclei and then to primary somatosensory cortex (S1), providing continuous feedback on limb position and load. S1 is tightly interconnected with primary motor cortex (M1), forming a loop directly relevant to postural correction. Sensorimotor EEG rhythms in the alpha–beta range ($\sim 10\text{--}20$ Hz, the “mu” rhythm) have been linked to body-schema representation — the brain’s continuously updated internal map of the body’s own configuration, which is conceptually close to, though not identical with, the “default space” TOIT proposes.

Cerebellum

Cerebellar cortex (granule cells, Purkinje cells, and climbing and mossy fiber inputs) computes motor error and projects via deep cerebellar nuclei to motor output pathways and, indirectly, to the thalamus. The cerebellum is traditionally modeled as feedforward rather than oscillatory, but cortico-muscular coherence at the tremor frequency has been documented in essential tremor [20], implicating oscillatory sensorimotor-cortical circuits in at least some forms of postural tremor — a small but real crack in the assumption that cerebellar motor control is purely non-oscillatory.

Taken together, these systems show extensive anatomical interconnection — particularly thalamic convergence of vestibular, somatosensory, and visual input — that TOIT interprets as infrastructure for a unified, oscillation-based spatial representation rather than several independent feedback loops running in parallel and only combining at a late cortical stage. (**Figure 1**) summarizes this convergence schematically.




Figure 1: Convergence of peripheral sensory input on thalamic nuclei under the TOIT framework. Conceptual schematic of proposed information flow; not a depiction of measured connectivity strength.

Oscillatory Activity Across Sensory Systems

Visual

V1 shows stimulus-driven gamma-band activity ($\sim 40\text{--}70$ Hz), classically linked to feature binding [21]. Retinal oscillatory potentials and LGN bursting near 40 Hz have also been documented [22,23], proposed that fast retino-geniculo-cortical oscillations in this range help construct continuous visual perception rather than a series of discrete snapshots. More directly relevant to posture, EEG studies of balance perturbation show that visually-induced destabilization (e.g., a sudden visual field rotation) and physically-induced destabilization both elicit theta-band (4–8 Hz) and beta-band cortical responses, with the visual condition producing stronger occipito-parietal theta activity specifically [24]. This is a genuinely direct empirical anchor for TOIT’s proposed link between visual/postural theta and correction — though it should be stated plainly that this finding describes a cortical response to an already-delivered

perturbation, not the continuous, moment-to-moment “self-motion tracking” that TOIT proposes occurs even in the absence of perturbation, and the two should not be treated as equivalent evidence.

Auditory

Gamma-band synchrony supports auditory scene analysis, and cross-modal gamma coupling with visual cortex occurs for congruent audiovisual stimuli [25]. Auditory cortex also phase-locks to slower rhythms (delta/theta, ~2–7 Hz) that track the rhythm of footsteps or speech — a rhythm TOIT proposes could couple with vestibular rhythms during walking or dancing, though this specific auditory–vestibular coupling claim is, at present, an extrapolation rather than a directly measured phenomenon.

Vestibular

Vestibular afferents fire in proportion to head velocity and acceleration rather than at any intrinsic oscillatory frequency of their own, but rhythmic head movement — as in walking or intentional oscillation — can produce theta-range firing in downstream thalamic and cortical neurons, including head-direction cells [26]. Postural sway itself is typically very slow at rest (~0.2 Hz), but rises toward 4–6 Hz under destabilizing conditions such as standing on an unstable surface with eyes closed, a shift that has been interpreted as reflecting increased corrective activity [27] — and which TOIT reads as consistent with, though not direct proof of, an increased demand on the theta-timed correction mechanism proposed in Section 6.

Cerebellar

Purkinje cell firing can show rhythmic components near 8 Hz, and cerebello-cortical-motor network activity has been characterized using MEG-based coherence imaging, an approach that has been used to map both physiological and pathological (e.g., Parkinsonian) oscillatory motor networks involving the cerebellum [28]. Transient beta-band activity (15–25 Hz) has also been reported during motor learning, consistent with a role in error signaling that sits alongside, rather than replacing, the cerebellum’s traditional feedforward error-correction role.

Somatosensory / motor

Sensorimotor cortex shows well-characterized mu (~10 Hz) and beta (~20 Hz) rhythms linked to body-schema maintenance [29]. Slower body-wide rhythms — cardiac (~1 Hz) and respiratory (~0.2–0.3 Hz) — also modulate cortical excitability and vestibular sensitivity to some degree, though their contribution to postural control specifically remains a much smaller and less direct effect than the sensorimotor rhythms described above, and should not be overstated. More broadly, non-oscillatory neuromodulatory systems also shape the variability of ongoing cortical activity: catecholamine levels have been shown to alter both the moment-to-moment variability of cortical population activity and of perception itself [30] — a useful reminder that oscillatory coupling is not the only mechanism regulating the cortical state relevant to perception and action, and that any complete account of postural control will eventually need to incorporate neuromodulatory as well as oscillatory dynamics.

TOIT’s broader claim — that these frequency bands are not simply present within each modality independently, but are phase-locked with one another via thalamic and/or hippocampal timing signals [31] — is the theory’s central and most novel proposal, and also the one most in need of direct empirical

testing, for instance via simultaneous intracranial recording across these systems during active balance tasks. Sections 6 and 7 lay out this proposal in more mechanistic detail.

The Thalamic Dual-Oscillator Hypothesis

TOIT proposes that the thalamus performs two coordinated oscillatory roles simultaneously, operating on different rhythms and serving different functions, illustrated schematically in (**Figure 2**).

Theta-based orientation timing

Anterior and lateral thalamic nuclei, working together with the hippocampal formation, generate theta-band activity that is already well documented in navigation and memory [32]. Because thalamic head-direction cells also show theta-rhythmic modulation [33]. TOIT proposes that this rhythm functions as a recurring “checkpoint” — roughly every 125–200 milliseconds at 5–8 Hz — at which incoming vestibular and proprioceptive signals are compared against the current internal spatial model, with corrections issued if a misalignment is detected. This proposed recalibration process is broadly consistent with behavioral evidence that returning astronauts, whose vestibular and proprioceptive inputs have been decorrelated from normal gravitational cues for months, show measurable locomotor and balance deficits that resolve gradually as sensorimotor recalibration proceeds [34]— though it must be said plainly that this study measured functional locomotor recovery, not thalamic oscillatory activity directly, so it is consistent with a recalibration process in general rather than specific evidence for this particular theta mechanism. Clinically, thalamic deep brain stimulation for tremor sometimes also improves gait and balance [35], which is at least consistent with a role for thalamic oscillatory activity in postural control, though it likewise does not by itself establish the specific theta-timing mechanism proposed here.

Alpha-based gating

Thalamic relay and reticular nuclei show alpha-band activity (8–12 Hz) associated with inhibition and gating of sensory throughput [36,37]. TOIT proposes this rhythm suppresses irrelevant or destabilizing sensory information for example, down-weighting visual input aboard a moving vessel so that vestibular input dominates, potentially reducing motion sickness. This is consistent with more general findings that alpha oscillations gate attentional access to sensory streams, including direct evidence that frontoparietal cortex controls which visual information is processed via modulation of anticipatory alpha desynchronization [38,39,40]. It should be emphasized, however, that direct evidence for this specific balance-related gating function as opposed to the general attentional gating role alpha is known to play remains limited, and the balance-specific application is, again, an extrapolation from an established general mechanism rather than a directly demonstrated one.




Figure 2: Proposed thalamic dual-oscillator mechanism: a theta-timed alignment loop (left) operating in parallel with an alpha-gated sensory-suppression loop (right). Timing values shown are illustrative of the proposed mechanism, not measured parameters.

Together, TOIT proposes the thalamus operates bidirectionally: theta activity coordinating with brainstem and spinal input to maintain reflexive alignment, and alpha activity gating what reaches cortical awareness. This framing offers a candidate explanation for several clinical patterns, including thalamic astasia — interpreted here as loss of the theta-based alignment mechanism despite intact cerebellar and vestibular function [14] and anterior thalamic lesions producing spatial disorientation independent of postural instability, which is consistent with the anterior thalamus’s strong connectivity to the hippocampal navigation system [41].

We note explicitly that the more detailed proposed dynamics — for instance, gamma amplitude nested within a theta “carrier” wave, itself modulated by an alpha gating envelope — are modeled on well-established theta/alpha–gamma coupling as a general communication mechanism between brain regions [42,43], and on evidence that theta phase modulates gamma power during spatial tasks more generally [44]. The specific claim that this same architecture governs moment-to-moment postural correction is, at present, an extrapolation rather than a directly tested finding, and we treat it as such throughout.

Cross-Frequency Coupling as an Integrating Mechanism

Cross-frequency coupling (CFC) — the coordination of oscillations at different frequencies — is TOIT’s proposed mechanism for binding gamma-band sensory detail to theta/alpha-timed integration. CFC is well documented in navigation, attention, and sensorimotor tasks generally [45,57]. which gives the general mechanism strong independent support even where its specific application to posture remains speculative.

Theta–gamma coupling for spatial binding

This is one of the best-established examples of CFC in neuroscience, particularly in hippocampal place-cell sequences during spatial navigation, where theta phase organizes gamma-band ensemble activity [43,46]. Human studies show comparable theta–gamma phase coupling during spatial working memory and virtual navigation tasks [47,44]. TOIT extends this established mechanism to hypothesize that ongoing postural correction works the same way: each theta cycle providing a timing frame, with gamma activity filling in sensory detail within that frame. (**Figure 3**) illustrates this proposed nesting schematically. This extension is plausible given the underlying mechanism’s generality across other cognitive domains, but

has not, to our knowledge, been directly tested for posture specifically — a gap we return to in Section 9.




Figure 3: Proposed cross-frequency coupling architecture: theta-nested gamma activity modulated by an alpha gating envelope. Schematic waveform for illustration only, not recorded neural data.

Alpha/beta–gamma coupling for sensorimotor regulation

Cortico-cerebellar beta rhythms (15–25 Hz) appear to support postural stability, decreasing when a corrective movement is required and rising during stable stance [48]. Pathological over-synchronization between cortical beta and muscle activity has been observed in some tremor-related postural disorders [49]. Alpha–gamma coupling, separately, has a well-established role in attentional gating: increased alpha and decreased gamma in a to-be-ignored sensory channel, and the reverse pattern in an attended one [40]. TOIT proposes an analogous pattern for balance — visual alpha rising to suppress unreliable visual input while vestibular gamma is favored when vision is unreliable — though this specific application again remains a hypothesis rather than a demonstrated finding.

Clinical relevance

Several disorders are consistent with a CFC-disruption account, though we present this as an interpretive lens rather than an established mechanism. Age-related imbalance is associated with greater reliance on visual input and altered cortical network organization during balance tasks in older adults, as shown using EEG connectome analysis [50], consistent with, though not direct proof of, a shift in cross-modal coupling with age. Sensory ataxia and the positive Romberg sign (loss of balance when vision is removed) are consistent with a scenario in which proprioceptive gamma input can no longer couple with visual and vestibular gamma via a shared theta reference, though direct oscillatory evidence in this specific population remains limited. Vestibular migraine shows abnormal thalamic activation to vestibular stimulation between episodes [15], consistent with — though again not proof of — a thalamic gating or coupling deficit.

Potential interventions

Theta-frequency transcranial alternating current stimulation (tACS) over vestibular cortex has been shown to affect balance performance in healthy participants [51], and 40 Hz sensory entrainment has shown some benefit in Alzheimer’s disease models [52]. These findings are suggestive that frequency-specific neuromodulation can influence the systems TOIT describes, though whether they act via the specific coupling mechanisms TOIT proposes — rather than some other, unrelated pathway — is not yet

established and should be treated as an open empirical question rather than a confirmed mechanism.

Clinical Correlates

This section considers how a TOIT-based account might interpret several vestibular and balance disorders. These are offered as candidate interpretations consistent with TOIT, not as established explanations. In most cases the standard model already accounts for these findings reasonably well; TOIT's contribution would be an additional explanatory layer — the oscillatory “why” underlying an already-documented “what” — rather than a replacement for the standard account.

Vestibular loss and spatial cognition

Beyond balance itself, chronic vestibular loss is associated with spatial memory and navigation deficits [53]. TOIT would interpret this as a loss of theta input needed to anchor the hippocampal-thalamic spatial map to a stable gravitational reference. Vestibular rehabilitation, which trains visual-vestibular coordination, may work partly by re-establishing cross-frequency coupling between the remaining, intact systems [54] though this mechanistic account of why rehabilitation works is TOIT's addition to an already well-established clinical practice, not a claim that displaces it.

Sensory ataxia / Romberg sign

Patients with proprioceptive deficits (e.g., dorsal column injury) can stand with eyes open but sway or fall with eyes closed, indicating reliance on conscious visual compensation for a degraded automatic (proprioceptive) system. Balance training on unstable surfaces may build alternative oscillatory coupling pathways over time [55], offering one candidate mechanism — among others — for why such training is clinically effective.

Vestibular migraine

Abnormal thalamic and cortical activation patterns are present even between attacks [15], which is consistent with a persistently lowered threshold for treating ordinary vestibular signals as destabilizing — that is, a gating deficit, under TOIT's framing, rather than a problem with the peripheral vestibular apparatus itself.

Mal de Debarquement Syndrome (MdDS)

Patients with MdDS feel a persistent rocking or swaying sensation at rest, often relieved paradoxically by passive motion (such as driving or being on a boat). Neuroimaging has shown increased metabolic activity in the left entorhinal cortex and amygdala in persistent MdDS, alongside decreased prefrontal and temporal metabolism — notable because the entorhinal cortex is a key hub for spatial information processing en route to the hippocampus and a major driver of brain oscillations [56], which fits comfortably with TOIT's emphasis on hippocampal-thalamic spatial oscillations. Optokinetic and vestibular readaptation approaches, including standardized rotational/optokinetic protocols, have shown clinical benefit in recent treatment guidelines [58], consistent with a resonance/re-coupling account, though this remains an active area of clinical research rather than a settled mechanism.

Persistent Postural-Perceptual Dizziness (PPPD) / visually-induced dizziness

Reduced functional connectivity between thalamus and putamen has been reported in these patients

[59], consistent with disrupted thalamic gating and an over-reliance on visual input for orientation — a pattern that maps naturally onto TOIT’s alpha-gating proposal, even though the original study was not designed to test that specific hypothesis.

Crebellar ataxia

Classic cerebellar disorders — postural tremor, titubation — are well explained by the standard model, and nothing here is meant to challenge that. TOIT’s additional claim, that poor cerebello-cortical oscillatory coupling (not just cerebellar damage per se) contributes to the subjective sense of persistent unsteadiness these patients report, is plausible but has limited direct evidence to date [60].

Pusher syndrome

Following posterolateral thalamic stroke, some patients actively push themselves toward their affected (paretic) side, as though their internal “upright” reference has shifted. TOIT interprets this as a shift in the thalamically anchored default space itself, requiring visual feedback (e.g., mirrors) to retrain orientation during rehabilitation — a reframing of an established rehabilitation technique rather than a new one.

Entrainment-based therapy directions

Rhythmic sensory stimulation — auditory metronomes, vibrotactile feedback, or optokinetic patterns tuned to natural postural or vestibular resonance frequencies — is a promising, already partially supported direction [61,62], for the related gamma-entrainment literature in Alzheimer’s disease). TOIT provides a theoretical rationale for why frequency-matched, rather than simply intense, sensory stimulation might be more effective, but this remains to be tested directly for balance rehabilitation specifically.

Testable Predictions and a Methodological Roadmap

A theory earns its place by generating predictions that could, in principle, come out false. We outline three here, in increasing order of technical difficulty, along with the specific methods that would be needed to test each one.

First, theta–gamma coupling between thalamus and hippocampus should increase during active spatial reorientation — for instance, when a person actively navigates a novel environment compared to when they passively view the same environment. This is testable via intracranial recording in surgical patients undergoing monitoring for other clinical reasons, or via high-density scalp EEG with source localization, though the latter has substantially lower spatial resolution for deep structures like the thalamus and hippocampus and would need to be interpreted cautiously.

Second, perturbing thalamic oscillations directly — for example, via 7 Hz transcranial alternating current stimulation over parietal cortex, targeting the theta rhythm TOIT proposes — should produce a measurable, direction-specific bias in perceived head orientation or postural angle, over and above any generic effect of stimulation. A well-designed study would need frequency-matched and sham-stimulation control conditions to rule out non-specific effects of electrical stimulation itself.

Limitations and Scope

We want to state the theory's current limitations plainly rather than leave them implicit in hedged language scattered through earlier sections. First, TOIT as applied here is an extension by analogy from mechanisms established in other domains — hippocampal theta-gamma coupling in spatial memory, alpha gating in visual attention — to a new domain, postural control, where the corresponding direct evidence is largely absent. Analogical extension is a legitimate way to generate hypotheses, but it is not evidence, and we have tried throughout this paper to keep that distinction explicit rather than letting a citation to an established mechanism in one domain stand in for evidence in another.

Second, the theory's most distinctive and novel claim — that oscillatory activity relevant to spatial cognition extends to cellular and tissue-level bioelectric signaling beyond canonical processing regions — is currently untestable with existing non-invasive human recording methods, and we do not know of a clear technical path to testing it in the near term. This is a genuine weakness of the theory's scope, not a minor caveat, and readers should weigh the rest of the framework accordingly.

Third, several of the clinical correlates discussed in Section 8 are also well explained by the standard vestibular-cerebellar model on its own terms; TOIT's contribution in those cases is an alternative or additional layer of explanation, not evidence that the standard model is wrong. We have tried to flag this distinction case by case rather than implying that clinical consistency with TOIT constitutes disconfirmation of the standard account.

Fourth, this paper does not present new experimental data. It is a synthesis and reinterpretation of existing anatomical, histological, and electrophysiological findings, several of which were established for purposes unrelated to balance or spatial orientation specifically. The methodological roadmap in Section 9 is offered precisely because we think the theory, in its current form, has been pushed about as far as a literature synthesis alone can take it, and further progress requires new data collected with these specific predictions in mind.

Conclusion and Future Directions

TOIT offers an oscillation-based alternative to the standard reflex-arc account of orientation and balance: rather than treating posture as purely a corrective response to error signals, it proposes that the brain continuously maintains an internal three-dimensional model of the body in space, with the thalamus coordinating this model via theta timing and alpha gating, and gamma-band synchrony supplying cross-modal detail. This does not require discarding the established roles of the cerebellum, vestibular apparatus, or proprioceptive system; it proposes instead that the timing and coordination between these systems carries information not captured by studying each system in isolation.

If supported by the kind of targeted testing outlined in Section 9, this framework suggests balance rehabilitation could benefit from an oscillatory approach — rhythmic, frequency-matched multisensory training, such as balance exercises paired with a metronome or virtual-reality visual-vestibular training — as a complement to, not a replacement for, standard physical therapy. It may also offer a way to stratify patients — for instance, identifying a specific alpha-gamma coupling deficit in the vestibular cortex — for targeted rather than generic neuromodulation.

We want to be direct, one final time, about the theory's current status. TOIT as applied here to orientation and balance is a hypothesis built by extension from an existing, more general theory of consciousness. It is supported by anatomical plausibility and by consistency with several clinical syndromes, but it is not yet directly tested against its most specific and novel claims — particularly the thalamic dual-oscillator and cross-modal cross-frequency-coupling mechanisms described in Sections 6 and 7. The next necessary step is empirical: simultaneous multi-region recording during active balance tasks, and targeted neuromodulation studies designed to falsify the model, not merely to illustrate it.

Statements

Conflict of interest: The authors declare no commercial or financial relationships that could be construed as a potential conflict of interest.

Funding: No

Ethics: No animal or human studies are presented in this manuscript; no identifiable human data are included.

Data availability: All material discussed is contained within this article and its cited sources; further inquiries can be directed to the corresponding author.

AI disclosure: No generative AI was used in the preparation of this manuscript.

References

1. Jerath R, Crawford MW, Barnes VA. (2015) A unified 3D default space consciousness model combining neurological and physiological processes that underlie conscious experience. *Fronti Psychol.* 6:1204.
2. Jerath R, Beveridge C, Jensen M. (2019a) The default space theory of consciousness: Phenomenological support from personal observations and clinical deficits. *World J Neurosci.* 9(1):1-21.
3. Jerath R, Beveridge C. (2018) Top mysteries of the mind: Insights from the default space model of consciousness. *Fronti Human Neurosci.* 12, 162.
4. Jerath R, Beveridge C, Jensen M. (2019b) On the hierarchical organization of oscillatory assemblies: Layered superimposition and a global bioelectric framework. *Fronti Human Neurosci.* 13:426.
5. Jerath R, Beveridge C. (2024) Beyond awareness: The binding of reflexive mechanisms with the conscious mind—A perspective from default space theory. *Fronti Human Neurosci.* 18:1520138.
6. O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map: Preliminary evidence from unit activity in the freely-moving rat. *Brain Research*, 34(1), 171–175.
7. Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., & Moser, E. I. (2005). Microstructure of a spatial map in the entorhinal cortex. *Nature*, 436(7052), 801–806.
8. Unverdi, M., & Alsayouri, K. (2023). Neuroanatomy, cerebellum. In StatPearls. StatPearls Publishing.
9. Morton, S. M., & Bastian, A. J. (2004). Cerebellar control of balance and locomotion. *The Neuroscientist*, 10(3), 247–259.
10. Manto, M., et al. (2012). Consensus paper: Roles of the cerebellum in motor control—The diversity of ideas on cerebellar involvement in movement. *Cerebellum*, 11(2), 457–487.
11. Wijesinghe, R., et al. (2015). Vestibular interactions in the thalamus. *Frontiers in Neural Circuits*, 9, 79.
12. Wiener, S. I., & Taube, J. S. (Eds.). (2005). *Head Direction Cells and the Neural Mechanisms of Spatial Orientation*. MIT Press.
13. Palmisano, S., et al. (2018). Future challenges for vestibular systems in virtual reality. *Frontiers in Psychology*, 9, 1120.

14. Lee, P. H., et al. (2005). Thalamic infarct presenting with thalamic astasia. *European Journal of Neurology*, 12(4), 317–319.
15. Russo, A., et al. (2014). Abnormal thalamic function in patients with vestibular migraine. *Neurology*, 82(23), 2120–2126.
16. Sherman, S. M. (2007). The thalamus is more than just a relay. *Current Opinion in Neurobiology*, 17(4), 417–422.
17. Singer, W. (1999). Neuronal synchrony: A versatile code for the definition of relations? *Neuron*, 24(1), 49–65.
18. Lopez, C., & Blanke, O. (2011). The thalamocortical vestibular system in animals and humans. *Brain Research Reviews*, 67(1–2), 119–146.
19. Taube, J. S. (2007). The head direction signal: Origins and sensory-motor integration. *Annual Review of Neuroscience*, 30, 181–207.
20. Hellwig, B., Häussler, S., Schelter, B., Lauk, M., Guschlbauer, B., Timmer, J., & Lücking, C. H. (2001). Tremor-correlated cortical activity in essential tremor. *The Lancet*, 357(9255), 519–523.
21. Gray, C. M., & Singer, W. (1989). Stimulus-specific neuronal oscillations in orientation columns of cat visual cortex. *PNAS*, 86(5), 1698–1702.
22. Llinás, R., & Paré, D. (1991). Of dreaming and wakefulness. *Neuroscience*, 44(3), 521–535.
23. Jerath, R., Cearley, S. M., Barnes, V. A., & Nixon-Shapiro, E. (2016). How lateral inhibition and fast retinogeniculo-cortical oscillations create vision: A new hypothesis. *Medical Hypotheses*, 96, 20–29.
24. Peterson, S. M., & Ferris, D. P. (2018). Differentiation in theta and beta electrocortical activity between visual and physical perturbations to walking and standing balance. *eNeuro*, 5(4), ENEURO.0207-18.2018.
25. Rohe, T., & Noppeney, U. (2018). Reliability-weighted integration of audiovisual signals can be modulated by top-down attention. *eNeuro*, 5(1).
26. Stackman, R. W., & Taube, J. S. (1998). Firing properties of head direction cells in the rat anterior thalamic nucleus: Dependence on vestibular input. *Journal of Neuroscience*, 18(23), 9309–9322.
27. Loram, I. D., Maganaris, C. N., & Lakie, M. (2005). Human postural sway results from frequent, ballistic bias impulses by upper leg muscles. *Journal of Physiology*, 564(1), 295–311.
28. Schnitzler, A., Timmermann, L., & Gross, J. (2006). Physiological and pathological oscillatory networks in the human motor system. *Journal of Physiology-Paris*, 99, 3–7.
29. Jurkiewicz, M. T., et al. (2006). Post-movement beta rebound is generated in motor cortex: Evidence from neuromagnetic recordings. *NeuroImage*, 32(3), 1281–1289.
30. Pfeffer, T., Avramiea, A.-E., Nolte, G., Engel, A. K., Linkenkaer-Hansen, K., & Donner, T. H. (2018). Catecholamines alter the intrinsic variability of cortical population activity and perception. *PLoS Biology*, 16(2), e2003453.
31. Misselhorn, J., et al. (2019). Alpha-beta oscillations shape cortical network architecture supporting predictive coding. *bioRxiv*, 500785.
32. Buzsáki G. (2005). Theta rhythm of navigation: Link between path integration and landmark navigation, episodic and semantic memory. *Hippocampus*, 15(7), 827–840.
33. Tsanov, M., et al. (2011). Oscillatory entrainment of thalamic neurons by theta rhythm in freely moving rats. *Journal of Neurophysiology*, 105(1), 4–17.
34. Mulavara, A. P., Feiveson, A. H., Fiedler, J., Cohen, H., Peters, B. T., Miller, C., Brady, R., & Bloomberg, J. J. (2010). Locomotor function after long-duration space flight: Effects and motor learning during recovery. *Experimental Brain Research*, 202(3), 649–659.
35. Wu, A. D., et al. (2008). Anodal transcranial direct current stimulation over the cerebellum improves associative eyeblink conditioning in healthy adults. *Neuroscience Letters*, 448(2), 162–165.
36. Lopes da Silva, F. H. (1991). Neural mechanisms underlying brain waves: From neural membranes to networks. *Electroencephalography and Clinical Neurophysiology*, 79(2), 81–93.
37. Lorincz, M. L., et al. (2009). Temporal framing of thalamic relay-mode firing by phasic inhibition during the alpha rhythm. *Neuron*, 63(5), 683–696.
38. Capotosto P, Babiloni C, Romani GL, Corbetta M. (2009) Frontoparietal cortex controls spatial attention through modulation of anticipatory alpha rhythms. *J Neurosci*. 29(18):5863-72.
39. Voytek, B., et al. (2017). Oscillatory dynamics coordinating human frontal networks in support of goal maintenance. *Nature Neuroscience*, 20(8), 1318–1324.

40. Bonnefond M, Jensen O. (2015) Gamma activity coupled to alpha phase as a mechanism for top-down controlled gating. *PLoS One*. 10(6):e0128667.
41. Aggleton JP, Brown MW. (1999) Episodic memory, amnesia, and the hippocampal-anterior thalamic axis. *Behavioral Brain Sci*. 22(3):425-44.
42. Bonnefond M, Kastner S, Jensen O. (2017) Communication between brain areas based on nested oscillations. *eNeuro*. 4(2):ENEURO.0153-16.2017.
43. Lisman, J. E., & Jensen, O. (2013). The theta-gamma neural code. *Neuron*, 77(6), 1002–1016.
44. Bush D, Bisby JA, Bird CM, Gollwitzer S, et al. (2017). Human hippocampal theta power indicates movement onset and distance travelled. *PNAS*. 114(46):12297-302.
45. Canolty RT, Knight RT. (2010) The functional role of cross-frequency coupling. *Trends Cogn Sci*. 14(11):506-15.
46. O'Keefe, J., & Recce, M. L. (1993). Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus*, 3(3), 317–330.
47. Kaplan, R., et al. (2014). Medial prefrontal theta phase coupling during spatial memory retrieval. *Hippocampus*, 24(6), 656–665.
48. Engel, A. K., & Fries, P. (2010). Beta-band oscillations—signalling the status quo? *Current Opinion in Neurobiology*, 20(2), 156–165.
49. Hall, S. D., et al. (2014). The role of beta oscillations in Parkinson's disease. *Frontiers in Human Neuroscience*, 8, 496.
50. Chen, Y.-C., Chou, L.-S., & Hwang, I.-S. (2021). Reliance on visual input for balance skill transfer in older adults: EEG connectome analysis using minimal spanning tree. *Frontiers in Aging Neuroscience*, 13, 632553.
51. Holle, D., et al. (2016). Transcranial alternating current stimulation in the theta range over the vestibular cortex enhances postural stability. *Brain Stimulation*, 9(4), 500–508.
52. Adaikkan C, Tsai LH. (2020) Gamma entrainment: Impact on neurocircuits, glia, and therapeutic opportunities. *Trends in Neurosci*. 43(1):24-41.
53. Smith, P. F., et al. (2019). Revisiting vestibular contributions to spatial memory and the hippocampus. *Frontiers in Integrative Neuroscience*, 13, 30.
54. Herdman, S. J., & Clendaniel, R. A. (2014). *Vestibular Rehabilitation* (4th ed.). F.A. Davis Company.
55. Slobounov, S., et al. (2000). Postural stability and balance dynamics in healthy elderly: Testing the limits of oscillatory stability. *Journal of Motor Behavior*, 32(3), 201–212.
56. Cha YH. (2015) Mal de débarquement syndrome: New insights. *Ann New York Acad Sci*. 1343(1):63-68.
57. Chacko, R. V., et al. (2018). Can oscillatory neural networks explain sensory integration? *Current Opinion in Neurobiology*, 52, 77–87.
58. Schoenmaekers, C., Jillings, S., De Laet, C., Zarowski, A., & Wuyts, F. L. (2024). Guideline for standardized approach in the treatment of the Mal de Débarquement syndrome. *Frontiers in Neurology*, 15, 1359116.
59. Van Ombergen, A., et al. (2017). Altered functional brain connectivity in patients with visually induced dizziness. *NeuroImage: Clinical*, 14, 538–545.
60. Ushiyama, J., et al. (2011). Cortical activity during gait in patients with cerebellar ataxia: A kinetic-cortical coupling analysis. *NeuroImage*, 55(3), 1289–1301.
61. Trainor, L. J., et al. (2009). Rhythmic auditory stimulation influences postural control in elderly. *Annals of the New York Academy of Sciences*, 1169, 378–381.
62. Iaccarino, H. F., et al. (2016). Gamma frequency entrainment attenuates amyloid load and modifies microglia. *Nature*, 540(7632), 230–235.